

Fungal Genetics Reports

Volume 24

Article 7

Cold-sensitive growth phenotype of some supersuppressor strains of *Neurospora crassa*

P. J. Russell
Reed College

Follow this and additional works at: <https://newprairiepress.org/fgr>



This work is licensed under a [Creative Commons Attribution-Share Alike 4.0 License](https://creativecommons.org/licenses/by-sa/4.0/).

Recommended Citation

Russell, P. J. (1977) "Cold-sensitive growth phenotype of some supersuppressor strains of *Neurospora crassa*," *Fungal Genetics Reports*: Vol. 24, Article 7. <https://doi.org/10.4148/1941-4765.1726>

This Research Note is brought to you for free and open access by New Prairie Press. It has been accepted for inclusion in Fungal Genetics Reports by an authorized administrator of New Prairie Press. For more information, please contact cads@k-state.edu.

Cold-sensitive growth phenotype of some supersuppressor strains of *Neurospora crassa*

Abstract

Cold-sensitive growth phenotype of some supersuppressor strains

Russell, P. J. Cold-sensitive growth phenotype of
some supersuppressor strains of *Neurospora crassa*.

The growth rates of a number of supersuppressor strains were determined in race tubes at 25 and 10°C to investigate whether any exhibited a cold-sensitive growth phenotype. Each race tube was inoculated with a conidial suspension and the lateral mycelial extension was marked at regular time intervals. All of the strains exhibited an approximately wild-type growth rate at 25°C. The steady-state growth rates for the supersuppressor strains and for the wild-type control at 10°C on both Vogel's complete and Vogel's minimal medium are presented in Table 1.

Table 1

Growth rates of supersuppressor strains and the wild-type in race tubes at 10°C

Strain	(FGSC#)	Growth rate (mm/hr)	
		Minimal	Complete
74A		0.68 (100)*	0.68 (100)*
ssu-1(?) a	(1750)	0.31 (46)	0.36 (53)
ssu-1;am A	(1687)	0.23 (34)	0.46 (68)
ssu-2;am A	(1689)	0.18 (27)	0.57 (84)
ssu-3;am a	(1851)	0.22 (33)	0.54 (80)
ssu-4;am A	(1852)	0.25 (37)	0.29 (43)
ssu-5 a	(1751)	0.32 (47)	0.42 (62)
ssu-6 A	(1748)	0.58 (85)	0.71 (104)

*% of wild-type rate on that medium.

At the gross observation level, it is significant to note that ssu-1(?) a on complete and ssu-3;am a on minimal showed a fairly regular "start-stop" type of growth, and strain ssu-2;am A exhibited very erratic growth characteristics on minimal. Thus the growth rates presented in Table 1 for these strains must be considered to be approximate.

If I arbitrarily consider a strain to be cold-sensitive if it exhibits a growth rate of 50% or less of that of the wild-type, then only ssu-4;am A is cold-sensitive on complete medium. By contrast, all of the strains except ssu-6 A are cold-sensitive on minimal medium. I conclude therefore, that the growth phenotype of these strains is medium-dependent. Nevertheless, it is interesting to note that most of the strains are cold-sensitive on the type of medium (minimal) that is used for testing the supersuppression phenotype (see, for example, Seale 1976 Mol. Gen. Genet. 148: 105).

My main interest in the cold-sensitive nature of these

strains stems from the possibility that their supersuppression phenotype is the result of altered ribosomes. Analysis of the ribosomes synthesized by the cold-sensitive strains on minimal medium at 10° C showed that the two subunits are made in wild-type proportions. Two-dimensional acrylamide gel electrophoresis experiments are currently being done to investigate whether altered ribosomal proteins can be detected in the strains. (This research was supported by NSF and NIH grants). - - - Department of Biology, Reed College, Portland, Oregon 97202.